arXiv:1110.2698v1 [nucl-th] 12 Oct 2011

The $O(2)$ model in polar coordinates at nonzero temperature

Martin Grahl[(a)], Elina Seel[(a)], Francesco Giacosa[(a)], and Dirk H. Rischke[(a,b)]
[(a)]*Institute for Theoretical Physics, Goethe University,*
Max-von-Laue-Str. 1, D–60438 Frankfurt am Main, Germany and
[(b)]*Frankfurt Institute for Advanced Studies, Goethe University,*
Ruth-Moufang-Str. 1, D–60438 Frankfurt am Main, Germany

We study restoration of spontaneously broken symmetry at nonzero temperature in the framework of the $O(2)$ model by using polar coordinates. We apply the CJT formalism to calculate the masses and the condensate in the double-bubble approximation, both with and without a term that explicitly breaks the $O(2)$ symmetry. We find that, in the case with explicitly broken symmetry, the mass of the angular degree of freedom becomes tachyonic above a temperature of about 300 MeV. Taking the term that explicitly breaks the symmetry to be infinitesimally small, we find that the Goldstone theorem is respected below the critical temperature. However, this limit cannot be performed for temperatures above the phase transition. We find that, no matter whether we break the symmetry explicitly or not, there is no region of temperature in which the radial and the angular degree of freedom become degenerate in mass. These results hold also when the mass of the radial mode is sent to infinity.

PACS numbers: 11.10.Wx,12.39.Fe,11.30.Rd

I. INTRODUCTION

The $O(N)$ model is of great importance for the theory of critical phenomena [1] and has been intensely studied in different theoretical frameworks [2–9]. Using a mass term with a 'wrong' sign, the $O(N)$ symmetry is spontaneously broken to $O(N-1)$ in the vacuum. At sufficiently high temperature T, the $O(N)$ symmetry is restored. The case $N=4$ has been profusely investigated because of its application to the chiral phase transition in quantum chromodynamics (QCD) [10–15]. QCD has a chiral $SU(N_f)_R \times SU(N_f)_L$ symmetry which is spontaneously broken to $SU(N_f)_{R+L}$ in the vacuum. The group $O(4)$ is locally isomorphic to $SU(2)\times SU(2)$, and $O(3)$ is locally isomorphic to $SU(2)$. Therefore, using the universality hypothesis, the $O(4)$ model can be considered as an effective theory for the restoration of chiral symmetry in QCD with two quark flavors.

Most studies of symmetry restoration within the $O(N)$ model [10–12, 14, 15] have been performed in Cartesian coordinates, $\vec{\phi} = (\phi_1, \phi_2, \ldots, \phi_N)$. Then, when the symmetry is spontaneously broken, the first coordinate is usually selected to assume a non-vanishing vacuum expectation value φ, $\phi_1 \to \varphi + \phi_1$, and the remaining $N-1$ coordinates are taken to be the $N-1$ Goldstone modes arising from spontaneous symmetry breaking. However, in the absence of explicit symmetry breaking terms, the effective potential is of the well-known 'Mexican hat'-type, with any state on the circle $|\vec{\phi}| = \varphi = const.$ being energetically degenerate with the vacuum state φ. Therefore, the Goldstone modes actually correspond to angular variables when moving along the circle $|\vec{\phi}| = const.$, and the massive degree of freedom corresponds to the radial degree of freedom when trying to 'climb' the rim of the hat.

This observation constitutes the main motivation for the present paper where we study symmetry restoration within the $O(N)$ model in polar coordinates. For the sake of simplicity, we shall concentrate on the case $N=2$, because the transition from Cartesian to polar coordinates is particularly simple in this case. Nevertheless, the conceptual issues are the same in this case as for $N=4$. In a certain sense, the case $N=2$ corresponds to QCD with one flavor in the absence of the axial anomaly. For $N_f = 1$, the (most simple) anomaly term would introduce a linear term in the massive degree of freedom, corresponding to an explicit symmetry breaking term. This is different from the two-flavor case, i.e., $N=4$, which corresponds to the case of maximal anomaly or maximal $U(1)_A$ breaking. Due to the similarities with the chiral symmetry of QCD and its breaking and restoration at nonzero T, in the following we shall also refer to the transition in the $O(2)$ model as 'chiral transition'. The order parameter for the transition, φ, will be called 'chiral condensate', and the two degrees of freedom (the 'chiral partners') are referred to as the σ

meson and the pion. The so-called 'chiral limit' is the case where explicit symmetry breaking terms are sent to zero, in which case the pion becomes a true Goldstone boson.

Another motivation for our work is that, if we send the mass of the radial degree of freedom (the σ meson) to infinity, we just obtain the leading-order Lagrangian of chiral perturbation theory (χPT) [16–18], with the Goldstone boson (the pion) as angular degree of freedom. In our simplified framework of the $O(2)$ model, it is then possible to perform a test of the validity of angular variables at nonzero T.

The transition from Cartesian to polar coordinates represents a change of the representation of the same theory. Obviously, physical quantities should be independent from the adopted representation. This fact is ensured by the so-called S-matrix equivalence theorem [19] which states that the elements of the S-matrix do not change when performing a transformation of the fields. However, for actual calculations one must use a certain approximation scheme. Then, the results obtained in one representation are not necessarily equal to those obtained in another representation [20]. In this work we shall explicitly show that, by using the CJT formalism in the double-bubble approximation [21], quantities computed in polar coordinates are actually quite different from those evaluated in the standard Cartesian coordinates.

Our results are the following: (i) for explicit symmetry breaking, the pion mass becomes tachyonic at high T, signalizing a breakdown of the CJT formalism. (ii) When the explicit symmetry breaking term is sent to zero (which, in analogy to the case $N = 4$, we refer to as the chiral limit), the Goldstone boson (pion) remains massless at each T in polar coordinates, thus satisfying the Goldstone theorem. This is contrary to Cartesian coordinates where a nonzero value of the pion mass is obtained as soon as the temperature is switched on, see e.g. Ref. [12]. However, due to singular terms $\sim 1/\varphi$ in the gap equations, and because the order parameter $\varphi = 0$ above the phase transition, the model becomes ill-defined. (iii) Both with and without explicit symmetry breaking, there is no region of temperature in which the chiral partners become degenerate in mass. (iv) Similar results hold also in the nonlinear limit, i.e., when the radial excitation becomes infinitely heavy.

The reason why polar coordinates are problematic at high T can be traced back to the decreasing value of the chiral condensate φ: in fact, in polar coordinates there are interaction terms proportional to inverse powers of φ, which render the application of the CJT formalism (or any other resummation scheme) problematic when φ is too small. In order to circumvent this problem one can perform a slightly different transformation to polar coordinates, in which the polar coordinates are not defined with respect to the origin of the Cartesian coordinates [22]. In this way a smooth limit from Cartesian to polar coordinates is realized, in which all the results of Cartesian coordinates can be reobtained. Interestingly, it is possible to investigate these issues also in the very simple situation of a free Lagrangian, see Sec. IV for details. In the Cartesian representation the results for thermodynamical quantities, such as the pressure, are exact in this case. The deviations of the results in the polar representation from the Cartesian one explicitly show the limitations of polar coordinates.

A further issue of polar coordinates is the fact that the Jacobian associated with the field transformation is not unity: an additional term emerges in the transformation of the interaction measure, which is potentially relevant in the context of quantum field theory. In order to investigate this point we study a different field representation in terms of polar variables, in which the Jacobian is indeed unity. The qualitative picture does not change and the same conceptual issue of exploding interaction terms in the high-temperature region exists also in this case.

The paper is organized as follows: In Sec. II we write the $O(2)$ model in terms of polar coordinates. In Sec. III we apply the CJT formalism and present the numerical results. The simple case of a free Lagrangian is discussed in Sec. IV. In Sec. V the alternative representation with unit Jacobian is discussed. Finally, we outline our conclusions and give an outlook in Sec. VI. Our units are $\hbar = c = k_B = 1$; the metric tensor is $g_{\mu\nu} = \mathrm{diag}\,(+,-,-,-)$.

II. THE $O(2)$ MODEL IN POLAR COORDINATES

A. Tree level, zero temperature

The $O(2)$ model in Cartesian coordinates $\vec{\phi} = (\phi_1, \phi_2)$, including an explicit symmetry breaking term $\sim H$, is described by the Lagrangian

$$\mathcal{L}_{cart} = \frac{1}{2}\partial_\mu \vec{\phi} \cdot \partial^\mu \vec{\phi} + \frac{m^2}{2}\vec{\phi} \cdot \vec{\phi} - \frac{\lambda}{2}(\vec{\phi} \cdot \vec{\phi})^2 + H\phi_1 \ . \tag{1}$$

As usual, we consider the shift $\phi_1 \to \phi_1 + \phi$, where ϕ is a constant. At zero temperature the minimization of the potential $V(\phi_1 = \phi, \phi_2 = 0)$ leads to the minimum $\phi = \varphi$ satisfying the following gap equation:

$$m^2\varphi - 2\lambda\varphi^3 + H = 0 \ . \tag{2}$$

When $H = 0$ the symmetry of the Lagrangian (1) under $O(2)$ transformations is exact. For $m^2 > 0$ and $H = 0$ the global minimum is realized for $\phi \equiv \varphi \neq 0$, i.e., the ground state breaks the $O(2)$ symmetry spontaneously. The vacuum expectation value φ is referred to as 'chiral condensate' in our model. In the vacuum, the numerical value for φ is chosen to be the pion decay constant, $\varphi \equiv f_\pi = 92.4$ MeV. When $H \neq 0$ an additional explicit breaking of chiral symmetry is realized.

By shifting the fields around their values at the global minimum, one obtains the zero-temperature tree-level masses m_1 and m_2 as

$$m_1^2 = -m^2 + 6\lambda f_\pi^2 \ , \quad m_2^2 = -m^2 + 2\lambda f_\pi^2 = \frac{H}{f_\pi} \ . \tag{3}$$

It is clear that $m_2 \to 0$ for $H \to 0$, i.e., this particle represents the Goldstone boson emerging from the spontaneous breaking of chiral symmetry.

We now introduce polar coordinates (σ, π) through the transformation

$$\phi_1 = \sigma \cos\frac{\pi}{\phi} \ , \ \phi_2 = \sigma \sin\frac{\pi}{\phi} \ , \tag{4}$$

leading to

$$\mathcal{L}_{pol} = \frac{1}{2}\partial_\mu\sigma\partial^\mu\sigma + \frac{1}{2}\frac{\sigma^2}{\phi^2}\partial_\mu\pi\partial^\mu\pi + \frac{m^2}{2}\sigma^2 - \frac{\lambda}{2}\sigma^4 + H\sigma\cos\frac{\pi}{\phi} \ . \tag{5}$$

Just as above, one shifts the field σ as $\sigma \to \sigma + \phi$. At zero temperature the minimization of the potential leads to the same Eq. (2) for the minimum $\phi = \varphi$. Also the zero-temperature tree-level masses m_σ and m_π coincide with the expressions of Eq. (3):

$$m_\sigma^2 = m_1^2 = -m^2 + 6\lambda f_\pi^2 \ , m_\pi^2 = m_2^2 = \frac{H}{f_\pi} \ . \tag{6}$$

In order to extract m_π^2, we have expanded the cosine in Eq. (5).

B. Mathematical issues using polar coordinates: The Jacobian and the integration intervals

Denoting $\pi/\phi \equiv \theta$ and taking into account that the Jacobian of the transformation in Eq. (4) is σ, the partition function can be rewritten as

$$Z = \oint_{-\infty}^{\infty} \mathcal{D}\phi_1\mathcal{D}\phi_2 \ \exp\left(-\int_0^{1/T} d\tau \int_V d\vec{x}\, \mathcal{L}_{E,cart}\right) = \oint_0^{\infty} \mathcal{D}\sigma\, \sigma \oint_0^{2\pi} \mathcal{D}\theta \ \exp\left(-\int_0^{1/T} d\tau \int_V d\vec{x}\, \mathcal{L}_{E,pol}\right) \ , \tag{7}$$

where periodic boundary conditions are understood: $\phi_1(\tau,\vec{x}) = \phi_1(\tau+1/T,\vec{x})$, etc. The suffix E means that the Lagrangians are considered in Euclidean space.

The r.h.s. of Eq. (7) describes the partition function in polar coordinates. Due to the fact that the Jacobian is not unity and that both fields σ and θ do not vary between $(-\infty,\infty)$, the question arises if we can apply the usual Feynman rules to the Lagrangian $\mathcal{L}_{pol}$.

One can rewrite the contribution of the Jacobian as

$$\oint_0^\infty \mathcal{D}\sigma\,\sigma \longrightarrow \oint_0^\infty \mathcal{D}\sigma\,\exp\left(-\Lambda^4\int_0^{1/T} d\tau\int_V d\vec{x}\,\ln\frac{\sigma}{\phi}\right) \tag{8}$$

where Λ is a UV cutoff. It is evident that this term induces a divergent contribution to the effective action requiring regularization. In the framework of dimensional regularization the contribution of the Jacobian vanishes in virtue of Veltman's rule [23], see also the explicit perturbative analyses performed in Refs. [22, 24]. In this work, however, we follow a different way. In the present section we neglect the contribution arising from the Jacobian. This is at first sight not legitimate because we do not work in dimensional regularization, but with different regularization schemes (trivial and counter-term regularizations, see Appendix A and Sec. III F). In Sec. V we introduce polar coordinates in a slightly modified manner which corresponds to a unitary Jacobian. We shall then *a posteriori* verify that the features described in the present section are qualitatively unchanged also when including the effects of the Jacobian.

We now turn to the extension of the range of integration over the fields σ and π. The possibility to extend the angular integration interval, $\int_0^{2\pi}\mathcal{D}\theta \to \int_{-\infty}^{\infty}\mathcal{D}\theta$, originates from the periodicity of the integrand. This point is subtle since it is in general not possible to split a path integral over the interval $I = I_1 \cup I_2$ into the sum of two path integrals over the intervals I_1 and I_2, respectively. For potentials of the form $U(\sigma)$ and for 2π-periodic potentials $U(\sigma,\theta)$ (as in the present case) one can show that extending the range of integration simply yields a countably infinite overall constant which can be absorbed into a normalization constant [25].

The extension in the σ direction from $(0,\infty)$ to $(-\infty,\infty)$ can be achieved with the help of a modified Heaviside step function defined in such a way that its contribution vanishes in dimensional regularization [25]. Note that in this work either the trivial or the counter-term regularizations are used. Although we do not expect a qualitative change of our results, we plan to perform a study using dimensional regularization for the counter terms.

C. Shift of the potential

In this section we show how to circumvent the problems arising from the fact that polar coordinates are ill-defined at the origin. Inspired by Ref. [22], we first shift the potential along the ϕ_1-axis by an arbitrary amount $v > 0$. In this way the global minimum realized at the critical temperature (and above) is not located at $\phi = 0$, but at $\phi = v$. After the shift, the Lagrangian (1) reads

$$\mathcal{L}_v = \frac{1}{2}\partial_\mu\phi_1\partial^\mu\phi_1 + \frac{1}{2}\partial_\mu\phi_2\partial^\mu\phi_2 + \frac{m^2}{2}(\phi_1 - v)^2 + \frac{m^2}{2}\phi_2^2 - \frac{\lambda}{2}\left[(\phi_1 - v)^2 + \phi_2^2\right]^2 + H(\phi_1 - v)\ . \tag{9}$$

When performing the transformation to polar coordinates, Eq. (4), we obtain

$$\mathcal{L}_v = \mathcal{L}_{pol} - m^2 v\,\sigma\,\cos\theta + \frac{m^2}{2}v^2 - \frac{1}{2}\lambda v(v - 2\sigma\cos\theta)\left(2\sigma^2 + v^2 - 2\sigma v\cos\theta\right) - H\,v\ . \tag{10}$$

Note that obviously $\mathcal{L}_{v=0} = \mathcal{L}_{pol}$ from Eq. (5), thus the study of Sec. II A can be regarded as a special case of this more general treatment. After performing a Taylor expansion of the trigonometric functions about $\theta = 0$ and shifting $\sigma \to \sigma + \phi$, we can easily determine the additional tree-level contributions to the masses and the interaction vertices.

III. RESULTS AT NONZERO T

In this section we present the results at nonzero temperature for the model described by the Lagrangian $\mathcal{L}_v$ for different cases.

A. $H \neq 0$, $v = 0$

In this case the system is described by the Lagrangian (5) in Sec. II A. The effective potential in the CJT formalism reads

$$V_{\text{eff}} = U(\phi) + \frac{1}{2} \sum_{i=\sigma,\pi} \int_k \left[\ln G_i^{-1}(k) + D_i^{-1}(k,\phi)\; G_i(k) - 1 \right] + V_2 \left[\phi, G_\sigma, G_\pi\right] \;, \tag{11}$$

where U denotes the classical potential and the inverse tree-level propagators read

$$D_\sigma^{-1}(k,\phi) = -k^2 + m_\sigma^2 = -k^2 - m^2 + 6\lambda\phi^2 \;,\; D_\pi^{-1}(k,\phi) = -k^2 + \frac{H}{\phi} \;. \tag{12}$$

At nonzero T the condensate and the masses become T-dependent functions

$$\varphi \to \varphi(T),\, m_\sigma \to M_\sigma(T),\; m_\pi \to M_\pi(T) \;, \tag{13}$$

and the dressed propagators are given by

$$G_\sigma^{-1}(k) \equiv -k^2 + M_\sigma^2 \;,\; G_\pi^{-1}(k) \equiv -Z^2 k^2 + M_\pi^2 \;, \tag{14}$$

where Z is a wave-function renormalization factor for the pion.

The term V_2 in Eq. (11) is the contribution of all 2PI vacuum graphs, ϕ denotes the connected 1-point function in the presence of a source, and G denotes the full connected 2-point function in the presence of the source. In general, V_2 consists of infinitely many diagrams, which prohibits an explicit calculation of V_2. In practice, one therefore has to restrict oneself to certain classes of diagrams. We shall use the so-called *Hartree approximation* where only the double-bubble diagrams are taken into account:

$$V_2 = \frac{3}{2}\lambda \left[\int_k G_\sigma(k)\right]^2 - \frac{H}{8\phi^3}\left[\int_k G_\pi(k)\right]^2 - \frac{1}{2\phi^2}\left[\int_q G_\sigma(q)\right]\left[\int_k k^2 G_\pi(k)\right] \;.$$

By extremizing the effective potential in Eq. (11) we obtain the following equations:

$$m^2\varphi - 2\lambda\varphi^3 + H = 6\lambda\varphi \int_k G_\sigma(k) - \frac{H}{2\varphi^2}\int_k G_\pi(k) + \frac{3H}{8\varphi^4}\left[\int_k G_\pi(k)\right]^2 + \frac{1}{\varphi^3}\left[\int_q G_\sigma(q)\right]\left[\int_k k^2 G_\pi(k)\right] \;, \tag{15}$$

$$M_\sigma^2 = -m^2 + 6\lambda\varphi^2 + 6\lambda\int_k G_\sigma(k) - \frac{1}{\varphi^2}\int_k k^2 G_\pi(k) \;, \tag{16}$$

$$M_\pi^2 = \frac{H}{\varphi} - \frac{H}{2\varphi^3}\int_k G_\pi(k) \;, \tag{17}$$

$$Z^2 = 1 + \frac{1}{\varphi^2}\int_k G_\sigma(k) \;, \tag{18}$$

where $\phi = \varphi$ denotes, in general, an extremum. At $T = 0$ the masses coincide with their tree-level values in Eq. (3) and $\varphi = f_\pi$. Furthermore,

$$H = m_\pi^2 \varphi \;, \tag{19}$$



FIG. 1: Upper row ($H \neq 0$), from left to right: masses, extrema of the effective potential, pressure (dashed: Cartesian, solid lines: polar, $v = 0$). Lower row ($H = 0$), from left to right: masses, extrema of the effective potential, pressure (dashed: Cartesian, solid lines: polar, $v = 0$). The dotted and dash-dotted lines in the middle panel of the upper row correspond to the condensate in the polar case with $v = 0.5 f_\pi$ and $v = 0.7325 f_\pi$, respectively.

$$\lambda = \frac{m_\sigma^2 - m_\pi^2}{4\varphi^2} \ , \tag{20}$$

$$m^2 = \frac{m_\sigma^2 - 3m_\pi^2}{2} \ . \tag{21}$$

The following numerical values are used at zero temperature: $m_\sigma = 600$ MeV, $m_\pi = 139.5$ MeV, and $f_\pi = 92.4$ MeV.

In this section we present the results using the so-called trivial regularization: the vacuum part of the integrals is simply set to zero, for details see Appendix A. For a discussion of the alternative counter-term regularization scheme we refer to Sec. III F.

The numerical results for the masses, the condensate, and pressure can be found in the upper row of Fig. 1. The results of polar coordinates are also compared to the standard results obtained in Cartesian coordinates. One observes that in general the results in polar coordinates differ substantially from those in Cartesian coordinates, except for the pressure at low temperatures, where the agreement between solid and dashed curves is seen to be quite good. Moreover, although the condensate decreases sharply at a temperature ~ 300 MeV, indicating the onset of chiral symmetry restoration, the pion mass does not become degenerate with the σ mass for high T. On the contrary, above a temperature $T_{max} \simeq 300.5$ MeV (where the curves in Fig. 1 terminate) it becomes tachyonic, signalizing a *breakdown* of the model for large T. This fact shows a limitation of the application of angular variables at nonzero temperature.

The effective potential is shown in Fig. 2 for different temperatures. There is a region around the origin where the effective potential is not defined since no real-valued solutions exist. Due to the singular terms with inverse powers of ϕ there is no extremum at the origin. At low temperature there is only a global minimum at a large value $\phi \equiv \Phi > 0$. At a certain temperature a second minimum and a maximum (both at smaller values of ϕ) occur. At $T^* \simeq 279.6$ MeV both are at the same level, indicating a first-order phase transition. Above this temperature, the global minimum Φ moves closer and closer to the origin, but never becomes zero. Above $T_{max} \simeq 300.5$ MeV no real solutions to the system of equations exist at the global minimum.

The reason why no real solution exists above $T_{\rm max}$ is due to the fact that the pion mass becomes imaginary, and therefore the system is not stable. In order to see this in more detail, we show the *function* $M_\pi(\phi)$ at $T = T_{max}$ in Fig. 3 (solid line). We observe that $M_\pi(\phi)$ has an imaginary part below a value $\phi_{turn} \simeq 15$ MeV. At ϕ_{turn} the imaginary part vanishes and M_π becomes a positive definite, real- (but multi-)valued function of ϕ. At $T = T_{max}$, the value at ϕ_{turn} coincides with M_π at the global minimum of the effective potential. The solid line above ϕ_{turn} is the solution for $M_\pi(\phi)$ which has to be used to plot the effective potential. Below T_{max} the point ϕ_{turn} is located such that the extremum $\varphi > \phi_{turn}$, i.e., real solutions at the global minimum Φ exist. With increasing temperature Φ approaches ϕ_{turn} and finally hits the ill-defined region for T_{max}.



FIG. 2: $H \neq 0$, $v = 0$. Effective potential minus its value at its global minimum $\phi = \Phi$ for temperatures $T = 278.91$ MeV and $T^* = 279.6$ MeV.



FIG. 3: $H \neq 0$, $v = 0$, $T = 300.5$ MeV $\simeq T_{max}$. Upper figure: Pion mass as function of the variable ϕ. Solid line: $M_\pi(\phi)$ when only the real solution exists, $\phi \geq \phi_{turn}$. Dashed line: $\mathrm{Re}M_\pi(\phi)$, dotted line: $\mathrm{Im}M_\pi(\phi)$. Lower figure: effective potential as function of the variable ϕ. The latter can be plotted only for $\phi \geq \phi_{turn}$, i.e., where real-valued solution for the pion mass exist. For $\phi < \phi_{turn}$ also the effective potential becomes a complex function.



FIG. 4: $H = 0$, $v = 0$. Effective potential minus its value at its global minimum for $T = 0$ (dotted), $T = 0.1$ MeV, $T = 0.999 f_\pi$, $T^* = 2.787 f_\pi$ (solid lines) and $T = 2.87 f_\pi$ (dotdashed).

B. $H = 0$, $v = 0$

In this section we study the case where the explicit symmetry breaking parameter is set to zero, $H = 0$. Equations (15)–(18) simplify to

$$m^2\varphi - 2\lambda\varphi^3 = 6\lambda\varphi \int_k G_\sigma(k) \ , \ M_\sigma^2 = -m^2 + 6\lambda\varphi^2 + 6\lambda \int_k G_\sigma(k) \ , \tag{22}$$

$$Z^2 = 1 + \frac{1}{\varphi^2} \int_k G_\sigma(k) \ , \ M_\pi^2 = 0 \ . \tag{23}$$

Here, we have used Eq. (A5) from Appendix A and the fact that $M_\pi^2 = 0$ to eliminate the pion tadpole term in Eq. (17). The numerical results for the masses, the condensate, and the pressure as functions of temperature are shown in the lower row of Fig. 1. Note that $M_\pi^2 = 0$ indicates that the Goldstone theorem is fulfilled at each T below the chiral phase transition. This is a property which does not hold in Cartesian coordinates, see for instance the lower left panel of Fig. 1. Unfortunately, $M_\pi^2 = 0$ also above the transition, indicating that the chiral partners do not become degenerate in the chirally restored phase where $\varphi = 0$.

The effective potential (relative to its value at its global minimum) for different temperatures is shown in Fig. 4. For temperatures in the transition region, one clearly observes the features of a first-order phase transition, i.e., three minima separated by two maxima.

C. $H \to 0$, $v = 0$

The chiral limit, i.e., $H \to 0$, does not exist, since for arbitrarily small, but nonzero values of H, the effective potential has no extremum near the origin $\varphi = 0$. We have confirmed this numerically by taking successively smaller values of H. The reason for this non-analyticity of the chiral limit are the inverse powers of φ in Eqs. (15)–(18). When setting $H = 0$ from the beginning, this problem does not exist because the singular terms are eliminated right away. Therefore, we conclude that the system cannot describe the chirally restored phase and is limited to the chirally broken phase. Again, polar coordinates have demonstrated a limited range of applicability. Interestingly, this problem does not exist in Cartesian coordinates where taking the limit $H \to 0$ is not problematic in the chirally restored phase.

D. $H \neq 0$, $v \neq 0$

As argued in Refs. [9, 22], the non-analyticity at the origin $\varphi = 0$ encountered in the previous subsections can be eliminated by introducing a non-vanishing value for the parameter v, i.e., shifting the vacuum expectation value to nonzero values $\varphi = v$. Now we have to consider the full Lagrangian (10). In the double-bubble approximation, the following equations are obtained:

$$\begin{aligned} 0 = {} & -m^2(\varphi - v) + 2\lambda(\varphi - v)^3 - H + 6\lambda(\varphi - v)\int_k G_\sigma(k) \\ & + \left[\frac{m^2 v - H}{2\varphi^2} + \lambda(\varphi - v)\left(\frac{v}{\varphi} + \frac{v^2}{\varphi^2}\right)\right]\int_k G_\pi(k) - \left(\frac{3}{8}\frac{m^2 v - H - 2v^3\lambda}{\varphi^4} + \frac{4v^2\lambda}{\varphi^3} - \frac{v\lambda}{4\varphi^2}\right)\left[\int_k G_\pi(k)\right]^2 \\ & + \frac{1}{\varphi^3}\left[\int_q G_\sigma(q)\right]\left[\int_k k^2 G_\pi(k)\right] - \left(-\frac{4v^2\lambda}{\varphi^3} + \frac{3v\lambda}{\varphi^2}\right)\left[\int_q G_\sigma(q)\right]\left[\int_k G_\pi(k)\right] , \end{aligned} \tag{24}$$

$$M_\sigma^2 = 6\lambda(\varphi - v)^2 - m^2 + 6\lambda\int_k G_\sigma(k) - \frac{1}{\varphi^2}\int_k k^2 G_\pi(k) + 2\lambda\,\frac{v}{\varphi}\left(3 - 2\,\frac{v}{\varphi}\right)\int_k G_\pi(k) , \tag{25}$$

$$\begin{aligned} M_\pi^2 = {} & \frac{H - m^2 v}{\varphi} + 2\lambda\frac{v}{\varphi}(\varphi - v)^2 - 2\lambda\frac{v}{\varphi}\left(2\frac{v}{\varphi} - 3\right)\int_k G_\sigma(k) \\ & - \left[\frac{H - m^2 v}{2\varphi^3} + \lambda\frac{v}{\varphi^3}(\varphi - v)^2 - 6\lambda\frac{v^2}{\varphi^2}\right]\int_k G_\pi(k) , \end{aligned} \tag{26}$$

$$Z^2 = 1 + \frac{1}{\varphi^2}\int_k G_\sigma(k) . \tag{27}$$

In the limit $v \to \infty$ the system of Eqs. (24)–(27) reduces to the system of equations for Cartesian coordinates obtained in Ref. [12] [see Eqs. (28a,b), (30a) for $N = 2$ with the replacement $\varphi \to \varphi - v$]. This is easily explained by the fact that the radial as well as the angular variable are defined relative to the origin so that for large ϕ we have $\sin\frac{\pi}{\phi} \simeq \frac{\pi}{\phi}$ and $\cos\frac{\pi}{\phi} \simeq 1$. Hence, in the limit $v \to \infty$ polar coordinates become Cartesian:

$$\phi_1 = \phi\cos\frac{\pi}{\phi} + \sigma\cos\frac{\pi}{\phi} \simeq \phi + \sigma \ , \ \ \phi_2 = \phi\sin\frac{\pi}{\phi} + \sigma\sin\frac{\pi}{\phi} \simeq \pi \ .$$

Solving the system of Eqs. (24)–(27) in the case $H \neq 0$ yields the condensate $\varphi - v$ shown in the middle panel of the upper row of Fig. 1 (dotted and dash-dotted lines). For values $v \lesssim 0.73 f_\pi$ (for instance $v = 0.5 f_\pi$, dotted line) the manner in which the condensate $\varphi(T)$ is multi-valued reminds of the behavior for a first-order phase transition. However, for this to be the case, there would have to be a third solution for $\varphi - v$ near the origin. This solution does not exist and therefore there is neither a first-order transition nor a restored phase. Above $v \gtrsim 0.73 f_\pi$ (dash-dotted line) the behavior of $\varphi(T)$ smoothly approaches the known Cartesian behavior, as expected.

E. The limit $m_\sigma \to \infty$

In this section we study the properties of the nonlinear sigma model [26] in polar coordinates at nonzero temperature. This investigation is also interesting in comparison to chiral perturbation theory at nonzero temperature, see also Ref. [27] and refs. therein.

The nonlinear sigma model is obtained by taking the limit $m_\sigma \to \infty$ and keeping the ratio λ/m^2 fixed. Inserting Eqs. (20) and (21) into Eq. (16) we obtain

$$M_\sigma^2 = \frac{3m_\pi^2}{2f_\pi^2}\left[f_\pi^2 - \varphi^2 - \int_k G_\sigma(k)\right] - \frac{1}{\varphi^2}\int\limits_k k^2 G_\pi(k) + \frac{m_\sigma^2}{2f_\pi^2}\left[3\varphi^2 - f_\pi^2 + 3\int\limits_k G_\sigma(k)\right] . \tag{28}$$

For $m_\sigma \to \infty$, the last term dominates. Therefore, if this term is positive definite, also $M_\sigma^2 \to \infty$. We have to demand that the last term is positive definite in order to have real-valued solutions for M_σ. Since $\int_k G_\sigma(k) \to 0$ for $M_\sigma \to \infty$, the positive-definiteness of the last term requires that

$$\varphi \geq \frac{f_\pi}{\sqrt{3}}. \tag{29}$$

This means that the range $\varphi < f_\pi/\sqrt{3}$ cannot be described, neither for $H \neq 0$ nor for $H \to 0$ (i.e., $m_\pi \to 0$). Thus, the model is not applicable in the chirally restored phase where we expect $\varphi \simeq 0$. This result shows that it is not possible to describe the phase transition in polar coordinates (in the framework of the CJT formalism) when taking the limit to the nonlinear sigma model.

F. Dependence on the regularization procedure

It is important to verify how the results change when a different regularization scheme is employed. In this section we regularize the divergent vacuum contributions of the tadpole diagrams by applying the counter-term scheme. The vacuum contributions of the thermal tadpole integral

$$\int \frac{d^3\vec{k}}{(2\pi)^3}\frac{1}{2\sqrt{\vec{k}^2 + M^2}} \tag{30}$$

is not neglected as in the trivial regularization scheme but is first rewritten using the residue theorem,

$$\int \frac{d^4k}{(2\pi)^4}\frac{1}{k^2 + M^2} . \tag{31}$$

This integral is regularized by introducing a renormalization scale μ [12]:

$$Q_\mu = \int \frac{d^4k}{(2\pi)^4}\left[\frac{1}{k^2+M^2} - \frac{1}{k^2+\mu^2} - \frac{\mu^2 - M^2}{(k^2+\mu^2)^2}\right] = \frac{1}{(4\pi)^2}\left[M^2 \ln\left(\frac{M^2}{\mu^2}\right) - M^2 + \mu^2\right] . \tag{32}$$

Figure 5 shows the temperature dependence of the condensate and the masses in the case $H \neq 0$, $v = 0$, applying the trivial regularization and the counter-term regularization prescription. We set $\mu = m_\sigma$ to satisfy the constraint $Z(T=0) = 1$ for the vacuum value of the pion wave-function renormalization factor.

As one can see, inclusion of the counter terms does not qualitatively change the results: it is not possible to avoid the pion from becoming tachyonic at high temperatures. This holds also when using the counter-term regularization in the chiral limit. Goldstone's theorem is fulfilled since the pions are massless in the symmetry broken phase. However, as with the trivial regularization scheme, no chirally restored phase exists for this system when the limit $H \to 0$ is taken.

IV. THE INSTRUCTIVE EXAMPLE OF THE FREE LAGRANGIAN

In order to explain the problems related to polar coordinates, in this section we consider a free Lagrangian in Cartesian coordinates:

$$\mathcal{L}_{free-cart} = \frac{1}{2}\partial_\mu\vec{\phi}\cdot\partial^\mu\vec{\phi} - \frac{m^2}{2}\,\vec{\phi}\cdot\vec{\phi} , \tag{33}$$



FIG. 5: The condensate (left panel) and the meson masses (right panel) as functions of temperature in the case $H \neq 0$, $v = 0$ in the trivial regularization (solid line) and the counter-term regularization scheme with $\mu = m_\sigma$ (dashed line).

where $m^2 > 0$. The pressure can be calculated exactly:

$$p_{free-cart} = -\frac{1}{2}\int\limits_k \ln D_\sigma^{-1} - \frac{1}{2}\int\limits_k \ln D_\pi^{-1} \ ,$$

with $D^{-1}_{\sigma,\pi}(k,\phi) = -k^2 + m^2_{\sigma,\pi} = -k^2 + m^2$. Explicitly:

$$p_{free-cart} = -\frac{T}{\pi^2}\int\limits_0^\infty dk k^2 \ln\left(1 - e^{-\frac{\sqrt{k^2+m^2}}{T}}\right) \ . \tag{34}$$

Let us now perform the transformation to polar coordinates. The Lagrangian

$$\mathcal{L}_{polar} = \frac{1}{2}\partial_\mu\sigma\partial^\mu\sigma + \frac{1}{2}\frac{\sigma^2}{\phi^2}\partial_\mu\pi\partial^\mu\pi - \frac{m^2}{2}\sigma^2 \tag{35}$$

follows directly from Eq. (5) by setting $\lambda = 0$ and replacing $m^2 \to -m^2$. Note that, although the original Cartesian Lagrangian (33) is that of a noninteracting system, the transformation to polar coordinates introduces a momentum-dependent four-particle interaction.

The gap equations are given by $M_\sigma^2 = m^2$, $Z^2 = \infty$, $M_\pi^2 = 0$. Because the tadpole $\int_k k^2 G_\pi(k)$ vanishes for $M_\pi = 0$, see Eq. (A5), the pressure in polar coordinates (and in double-bubble approximation) has no contribution from the interaction term $\frac{1}{2}\frac{\sigma^2}{\phi^2}\partial_\mu\pi\partial^\mu\pi$ and is given by

$$p_{polar} = -\frac{T}{2\pi^2}\int\limits_0^\infty dk k^2 \ln\left(1 - e^{-\frac{\sqrt{k^2+m^2}}{T}}\right) + \frac{T^4\pi^2}{90} \ . \tag{36}$$

The pressure $p_{free-polar}$ represents the sum of one particle with mass m and one particle with mass $m = 0$. This latter contribution is, however, not correct and overestimates the exact result of Eq. (34). This simple example shows once more that care is needed when polar coordinates (and a certain many-body approximation) are used to study properties of a system at nonzero T.

V. AN ALTERNATIVE WAY TO INTRODUCE POLAR COORDINATES

A. The transformation

As we have discussed in Sec. II B, the Jacobian associated with the transformation to polar coordinates introduced in Eq. (4) is not unity. In this section we present an alternative polar representation (ψ, π) for the $O(2)$ linear σ-model, which is defined as:

$$\phi_1 = \sqrt{2\phi\psi}\cos\frac{\pi}{2\phi} \ , \ \phi_2 = \sqrt{2\phi\psi}\sin\frac{\pi}{2\phi} \ . \tag{37}$$

In this case the associated Jacobian remains unity. Here the massive scalar field acquiring a nonvanishing vacuum expectation value in case of spontaneously broken symmetry is represented by ψ.

The Lagrangian (1) expressed in terms of the fields ψ and π introduced in Eq. (37) reads

$$\begin{aligned}\mathcal{L} = {} & \frac{1}{2}(\partial_\mu\psi)^2 + \frac{1}{2}(\partial_\mu\pi)^2 - 2m^2\phi^2 - 8\lambda\phi^4 + 2H\phi - \frac{\psi^2}{2}\left(16\lambda\phi^2 + \frac{H}{2\phi}\right) - \frac{\pi^2}{2}\frac{H}{2\phi} \\ & - (\partial_\mu\psi)^2\frac{\psi}{2\phi} + (\partial_\mu\pi)^2\frac{\psi}{2\phi} - \frac{H}{8\phi^2}\psi\pi^2 + \frac{H}{8\phi^2}\psi^3 + (\partial_\mu\psi)^2\frac{\psi^2}{2\phi^2} - \frac{5H}{64\phi^3}\psi^4 + \frac{H}{192\phi^3}\pi^4 + \frac{H}{32\phi^3}\psi^2\pi^2 \\ & + \text{higher-order terms.}\end{aligned} \tag{38}$$

where the shift $\psi \to \psi + \phi$ has also been performed.

The corresponding inverse tree-level propagators and tree-level masses are given by:

$$\begin{aligned} D_i^{-1}(k;\phi) &= -k^2 + m_i^2,\ i = \psi, \pi \\ m_\psi^2 &= 16\lambda\phi^2 + \frac{H}{2\phi},\ m_\pi^2 = \frac{H}{2\phi}\ .\end{aligned} \tag{39}$$

Applying the CJT formalism we derive the effective potential within the double-bubble approximation:

$$V_{eff}(\phi; G_\pi, G_\psi) = 2m^2\phi^2 + 8\lambda\phi^4 - 2H\phi + \frac{1}{2}\int[\ln G_\psi^{-1}(k) + D_\psi^{-1}(k;\phi)G_\psi(k) - 1] \tag{40}$$

$$+ \frac{1}{2}\int[\ln G_\pi^{-1}(k) + D_\pi^{-1}(k;\phi)G_\pi(k) - 1] + V_2(\phi; G_\pi, G_\psi)\ , \tag{41}$$

$$\begin{aligned} V_2(\phi; G_\pi, G_\psi) = {} & -\frac{1}{2\phi^2}\int_k k^2 G_\psi(k)\int_l G_\psi(l) + \frac{15H}{64\phi^3}\left[\int_k G_\psi(k)\right]^2 \\ & - \frac{H}{64\phi^3}\left[\int_k G_\pi(k)\right]^2 - \frac{H}{32\phi^3}\int_k G_\psi(k)\int_k G_\pi(k)\ .\end{aligned} \tag{42}$$

Finally, the stationary conditions for the effective potential give the following gap equations for the temperature-dependent masses, the condensate and the wave-function renormalization for the ψ-field.

$$\begin{aligned} 2H = {} & 4m^2\varphi + 32\lambda\varphi^3 + \left(16\lambda\varphi - \frac{H}{4\varphi^2}\right)\int_k G_\psi(k) - \frac{H}{4\varphi^2}\int_k G_\pi(k) + \frac{1}{\varphi^3}\int_k k^2 G_\psi(k)\int_l G_\psi(l) \\ & - \frac{45H}{64\varphi^4}\left[\int_k G_\psi(k)\right]^2 + \frac{3H}{64\varphi^4}\left[\int_k G_\pi(k)\right]^2 + \frac{3H}{32\varphi^4}\int_k G_\psi(k)\int_k G_\pi(k)\ ,\end{aligned} \tag{43}$$

$$M_\psi^2 = 16\lambda\varphi^2 + \frac{H}{2\varphi} + \frac{H}{16\varphi^3}\left[15\int_k G_\psi(k) - \int_k G_\pi(k)\right] - \frac{1}{\varphi^2}\int_k k^2 G_\psi(k)\ , \tag{44}$$

$$M_\pi^2 = \frac{H}{2\varphi} - \frac{H}{16\varphi^3}\left[\int_k G_\psi(k) + \int_k G_\pi(k)\right]\ ,\quad Z_\psi^2 = 1 + \frac{1}{\varphi^2}\int_k G_\psi(k)\ . \tag{45}$$

B. Results

In this subsection we present the numerical results for this alternative polar representation. Figure 6 shows the condensate and the masses as function of T using the trivial and the counter-term regularization schemes in the case of explicit chiral symmetry breaking. In Fig. 7 the same quantities are shown in the chiral limit. In the case of explicitly broken symmetry there is a crossover transition and one observes no degeneration of the chiral partners at high T. Just as in Sec. III, for $T \gtrsim 300$ MeV the validity of the model breaks down and the numerical results are no longer reliable. Moreover, chiral restoration is approached very slowly, also when the chiral limit is taken.



FIG. 6: The temperature-dependent condensate (left panel) and the meson masses (right panel) for the alternative polar representation in the case of explicitly broken symmetry, in the trivial regularization (solid lines) and in the counter-term regularization scheme (dashed lines) with $\mu = m_\psi$.



FIG. 7: The temperature-dependent condensate and ψ-mass for the alternative polar representation in the chiral limit, in the trivial regularization (solid line) and in the counter-term regularization scheme (dashed line) with $\mu = m_\psi$.

Summarizing the results the alternative polar representation does not offer a better description of the chiral phase transition (at least not within the chosen many-body approximation). The reason for this is that, when constructing this alternative polar representation, one has to evaluate several terms of the form $f(1+x)$ with $x = \psi/\phi$ and $x = \pi/\phi$. Increasing the temperature the condensate starts melting while the fluctuation of the fields become larger and the calculation becomes no longer reliable for higher temperatures. Especially when approaching the critical temperature all the terms of the form x/ϕ, with $x = \pi, \psi$, become problematic, limiting the validity of the model to lower temperatures.

VI. CONCLUSIONS

In this paper, we have studied the $O(2)$ model in polar coordinates at nonzero temperature. After having clarified some issues related to the transformation from Cartesian to polar coordinates in the functional integral representation of the partition function, we have computed the latter in the CJT formalism in double-bubble approximation. We have studied in detail the cases where the chiral symmetry is explicitly broken and where the explicit symmetry breaking parameter is set to zero. We have distinguished the latter case from the chiral limit, where the explicit symmetry breaking parameter is smoothly sent to zero. We have found that this limit does not exist in the strict mathematical sense, due to non-analytic terms

$$\frac{1}{2}\frac{\sigma^2}{\phi^2}\partial_\mu\pi\partial^\mu\pi \quad \text{and} \quad H\sigma\cos\frac{\pi}{\phi}$$

appearing in the Lagrangian for polar coordinates. For all cases except when the explicit symmetry breaking parameter is exactly zero, we have found that, when approaching $\phi \to 0$, the ensuing divergences are sufficiently severe to invalidate the approach above a certain maximal temperature T_{max}. Above T_{max} no physical solutions exist. The same results hold in the nonlinear limit $m_\sigma \to \infty$.

We have also investigated the possibility of shifting the potential by an amount v along the ϕ_1-direction in order to circumvent the divergences resulting from the non-analytic terms in the Lagrangian. Variation of v allows to change smoothly from polar coordinates to Cartesian coordinates (corresponding to $v \to \infty$). Our conclusions about the suitability of polar coordinates remain unchanged, however, T_{max} increases with v, so that the range of applicability of polar coordinates is extended. Morevoer, the larger v, the better the agreement with the Cartesian results.

We have also introduced and studied an alternative representation for polar coordinates (with unit Jacobian in the functional integral representation of the partition function). Also in this case, the above mentioned problems persist.

Our most important result is that, both in the chiral limit and in the case of explicitly broken symmetry, the chiral partners do not become degenerate in mass at high T, not even when approaching T_{max} where the order parameter has already decreased by a substantial amont. In general, the sigma particle becomes more massive while the pion mass decreases or remains zero (in the absence of explicit symmetry breaking). Above T_{max} the pion even becomes tachyonic. The absence of degeneracy of the chiral partners means that an important indication for the restoration of chiral symmetry is missing when using polar coordinates. We conclude that the use of angular variables is not well suited for the study of the chiral phase transition.

Note that we used either trivial regularization or the counter-term renormalization scheme when treating the divergent vacuum contributions to the thermal integrals. Our results are qualitatively independent from the chosen regularization scheme. Due to consistency with the extension of the range of functional integration in polar coordinates, one should perform the renormalization procedure in dimensional regularization, which remains a task for the future.

A possible extension of the present study would be to use four-dimensional polar coordinates, corresponding to the $O(4)$ model. The polar $O(4)$ model has the advantage that three degrees of freedom can be identified with the three pions, π^0 and $\pi^\pm$, and the remaining one with their chiral partner, the sigma particle. The number of angular degrees of freedom could affect the behavior of the equations in the limit $\varphi \to 0$. However, we believe that this generalization will not fix the problem encountered in the polar version of the $O(2)$ model, since the divergences pointed out above are general.

Finally, although on a conceptual level each representation is equivalent, on a practical level the use of Cartesian coordinates is favorable to study thermodynamical properties of systems described by the $O(N)$ model. In connection to QCD, one should extend the model by incorporating all the relevant low-energy mesons: besides scalar and pseudoscalar particles, also vector and axial-vector degrees of freedom should be included in an enlarged $U(N)\times U(N)$ symmetry for a more realistic treatment of properties of QCD at nonzero T [28–31].

Acknowledgment: The authors thank Stefan Strüber for useful discussions. M.G. and E.S. thank HGS-HIRe for FAIR for funding.

Appendix A: Thermal integrals

We list the standard thermal integrals which were used for numerical calculations. Carrying out the Matsubara summation of the thermal tadpole integral gives

$$\int_k G(k) = \int \frac{d^3\vec{k}}{(2\pi)^3}\frac{1}{\sqrt{\vec{k}^2+M^2}}\left[\frac{1}{2}+\frac{1}{\exp\left(\sqrt{\vec{k}^2+M^2}/T\right)-1}\right] , \tag{A1}$$

consisting of a finite contribution,

$$Q_T(M) = \int \frac{d^3\vec{k}}{(2\pi)^3} \frac{1}{\sqrt{\vec{k}^2+M^2}} \frac{1}{e^{\sqrt{\vec{k}^2+M^2}/T}-1} = \int\limits_0^\infty \frac{dk}{2\pi^2}\, \frac{k^2}{\sqrt{k^2+M^2}} \frac{1}{e^{\sqrt{k^2+M^2}/T}-1} \ ,$$

and a divergent vacuum contribution,

$$Q_\mu(M) = \int \frac{d^3\vec{k}}{(2\pi)^3} \frac{1}{2\sqrt{\vec{k}^2+M^2}} = \int \frac{d^4k}{(2\pi)^4} \frac{1}{k^2+M^2} \ . \tag{A2}$$

For the explicit calculation we need the following integrals:

$$\int\limits_k G_\sigma = Q_\mu(M_\sigma) + Q_T(M_\sigma) \ , \tag{A3}$$

$$\int\limits_k G_\pi = \frac{1}{Z^2}\left[Q_\mu\left(\frac{M_\pi}{Z}\right) + Q_T\left(\frac{M_\pi}{Z}\right)\right] \ , \tag{A4}$$

$$\int\limits_k k^2 G_\pi = \frac{M_\pi^2}{Z^4}\left[Q_\mu\left(\frac{M_\pi}{Z}\right) + Q_T\left(\frac{M_\pi}{Z}\right)\right] \ . \tag{A5}$$

For the effective potential we need in addition:

$$\begin{aligned}\frac{1}{2}\int\limits_k \left[D_\sigma^{-1}G_\sigma - 1\right] &= \frac{1}{2}\int\limits_k \left[(-k^2+m_\sigma^2)\frac{1}{-k^2+M_\sigma^2} - \frac{-k^2+M_\sigma^2}{-k^2+M_\sigma^2}\right] \\ &= \frac{1}{2}(m_\sigma^2 - M_\sigma^2)\int\limits_k \frac{1}{-k^2+M_\sigma^2} = \frac{1}{2}(m_\sigma^2 - M_\sigma^2)\left[Q_\mu(M_\sigma) + Q_T(M_\sigma)\right] \ , \end{aligned} \tag{A6}$$

$$\begin{aligned}\frac{1}{2}\int\limits_k \left[D_\pi^{-1}G_\pi - 1\right] &= \frac{1}{2}\int\limits_k \left[(-k^2+m_\pi^2)\frac{1}{-Z^2k^2+M_\pi^2} - \frac{-Z^2k^2+M_\pi^2}{-Z^2k^2+M_\pi^2}\right] \\ = \frac{1}{2}(m_\pi^2 - M_\pi^2)\frac{1}{Z^2}&\left[Q_\mu\left(\frac{M_\pi}{Z}\right) + Q_T\left(\frac{M_\pi}{Z}\right)\right] + \frac{1}{2}(Z^2-1)\frac{M_\pi^2}{Z^4}\left[Q_\mu\left(\frac{M_\pi}{Z}\right) + Q_T\left(\frac{M_\pi}{Z}\right)\right] \ , \end{aligned} \tag{A7}$$

$$\frac{1}{2}\int\limits_k \ln G_\sigma^{-1} = \frac{1}{2}R_\mu(M_\sigma) + \frac{1}{2}R_T(M_\sigma) \ , \tag{A8}$$

$$\frac{1}{2}\int\limits_k \ln G_\pi^{-1} = \frac{1}{2}R_\mu\left(\frac{M_\pi}{Z}\right) + \frac{1}{2}R_T\left(\frac{M_\pi}{Z}\right) \ , \tag{A9}$$

where

$$R_T\left(\frac{M}{Z}\right) = \frac{T}{\pi^2}\int\limits_0^\infty dk k^2 \ln\left(1 - e^{-\sqrt{k^2+\frac{M^2}{Z^2}}/T}\right) \ , \tag{A10}$$

which in the case $M = 0$ and $Z \neq 0$ simplifies to

$$R_T(0) = -\frac{T^4\pi^2}{45} \ . \tag{A11}$$

Except for the discussion in Secs. III F and V, we neglected the contributions from renormalization, i.e., $Q_\mu \equiv 0$ as well as $R_\mu \equiv 0$. This approximation scheme is called the trivial regularization.

---

[1] H. Kleinert and V. Schulte-Frohlinde. Critical properties of phi**4-theories. River Edge, USA: World Scientific (2001) 489 p.

[2] H. Meyers-Ortmanns, H.J. Pirner, and B.J. Schaefer. *Phys.Lett.*, B311:213–218, 1993.

[3] Heui-Seol Roh and T. Matsui. *Eur.Phys.J.*, A1:205–220, 1998.

[4] Jurgen Baacke and Stefan Michalski. *Phys.Rev.*, D67:085006, 2003.

[5] Jens O. Andersen, Daniel Boer, and Harmen J. Warringa. *Phys.Rev.*, D70:116007, 2004.

[6] Nicholas Petropoulos. *hep-ph/0402136*, 2004. Based on Ph.D. thesis submitted to Manchester University in Oct. 2000.

[7] Yu.B. Ivanov, F. Riek, and Joern Knoll. *Phys.Rev.*, D71:105016, 2005.

[8] Bao-Chun Li and Mei Huang. *Phys.Rev.*, D80:034023, 2009.

[9] E.N. Argyres, M.T.M. van Kessel, and R.H.P. Kleiss. *Eur.Phys.J.*, C64:319–349, 2009.

[10] Robert D. Pisarski and Frank Wilczek. *Phys. Rev.*, D29:338–341, 1984.

[11] Frank Wilczek. *Int. J. Mod. Phys.*, A7:3911–3925, 1992.

[12] Jonathan T. Lenaghan and Dirk H. Rischke. *J. Phys.*, G26:431–450, 2000.

[13] Agostino Butti, Andrea Pelissetto, and Ettore Vicari. *JHEP*, 08:029, 2003.

[14] H. Meyer-Ortmanns and T. Reisz. Principles of phase structures in particle physics. Hackensack, USA: World Scientific (2007) 682 p.

[15] Elina Seel, Stefan Struber, Francesco Giacosa, and Dirk H. Rischke. Available from arXiv:1108.1918.

[16] J. Gasser and H. Leutwyler. *Annals Phys.*, 158:142, 1984.

[17] G. Ecker, J. Gasser, A. Pich, and E. de Rafael. *Nucl.Phys.*, B321:311, 1989.

[18] Stefan Scherer. *Adv.Nucl.Phys.*, 27:277, 2003. To be edited by J.W. Negele and E. Vogt.

[19] S. Kamefuchi, L. O'Raifeartaigh, and Abdus Salam. *Nucl.Phys.*, 28:529–549, 1961.

[20] S. Kondratyuk, A.D. Lahiff, and H.W. Fearing. *Phys.Lett.*, B521:204–210, 2001.

[21] John M. Cornwall, R. Jackiw, and E. Tomboulis. *Phys. Rev.*, D10:2428–2445, 1974.

[22] E. N. Argyres, C. G. Papadopoulos, M. T. M. van Kessel, and R. H. P. Kleiss. *Eur. Phys. J.*, C61:495–518, 2009.

[23] George Leibbrandt. *Rev. Mod. Phys.*, 47:849, 1975.

[24] H. Kleinert and A. Chervyakov. *Eur.Phys.J.*, C19:743–747, 2001.

[25] Martin Grahl. *The $O(N = 2)$ model in polar coordinates at nonzero temperature.* Diploma thesis, Goethe-University Frankfurt am Main, 2009.

[26] Alexander Bochkarev and Joseph I. Kapusta. *Phys.Rev.*, D54:4066–4079, 1996.

[27] Stefan Leupold. Available from hep-ph/0407197.

[28] S. Gasiorowicz and D.A. Geffen. *Rev.Mod.Phys.*, 41:531–573, 1969.

[29] Pyungwon Ko and Serge Rudaz. *Phys.Rev.*, D50:6877–6894, 1994.

[30] Stefan Struber and Dirk H. Rischke. *Phys.Rev.*, D77:085004, 2008.

[31] Denis Parganlija, Francesco Giacosa, and Dirk H. Rischke. *Phys.Rev.*, D82:054024, 2010.